GENETICS

A PERIODICAL RECORD OF INVESTIGATIONS
BEARING ON HEREDITY AND
VARIATION

VOLUME 16 - 1931

WITH ELEVEN PLATES AND ONE HUNDRED TWENTY-FIVE TEXT FIGURES

PUBLISHED BY THE
BROOKLYN BOTANIC GARDEN
BROOKLYN, NEW YORK
U. S. A.

INDEX

Page numbers printed in black-faced type refer to headings of articles or of sections. Names in lists of "Literature cited" are not included.

AASE, H. C., 310, 446
Aceto-carmine
  Method, 327
  Mount, 201, 477
  Smear, 597
Aegilops, 446
  cylindrica, 446
  Hybrids, 446
  ovata, 447
  speltoides, 446
  triuncialis, 447
Agar, 228
AGOL, I. J., 254
Algae, 587
  Blue green, 144
  Dormancy, 587
Allelomorphs, multiple, 104
ALLEN, G. M., 45, 50, 69, 160
Allen's modification of Bouin's fixative, 424, 477
ALMON, L., 624
ALTENBURG, E., 490, 630, 637
ALTSHULER, V. E., 254
All-Union Communistic Academy, 254
All-Union Meeting of Genetics and Selection, 254
American Association for the Advancement of Science, 133, 659
Anaphylactic shock in pigeons, 412
ANDERSON, E. G., 375, 376, 379, 381, **386**, 387, 389, 392, 393, 602, 625
Aneuploid forms in Fragaria, **455**, 464
Angiosperms, 589
  Dormancy, 589
  Vivipary, 589
Antisera, rodent, 398, 406, 418
Aphids, 251
Apotettix, **353**–355, 374
  eurycephalus, 353
Artemisia nitida, 478
Assortative mating, 146
Assortment, random, 146
Autosomal genes, **364**, **368**, 371, 373
Autosomal linkage, 373
Avena barbata, 478
Avena strigosa, 478
AVERY, P., 445, 450
Avicenna, 589
AZEVEDO, J. L., 415, 416

BABCOCK, E. B., 323
Backcrossing, 191–198, **204**, 205, 208, 210
BACKUS, M. L., 624
Bacteria, 144
BAGG, H. J., 54, 55
BAILEY, H. W., 289
Banana agar, 282
Bantams, Sebright, 560
Barley, 291, 300, 529
BARNES, H. F., 252
BARTLETT, H. H., 345, 346
BATESON, W., 50
BAUR, E., frontispiece, xi
BEADLE, G. W., 190, 479
Beans, 5, 6
  Effect of cross-pollination on the size of seed, 5
BEHRENS, 402
BELAR, K., 251
BELLING, J., 92, 342, 344, 346, 347, 460, 479, 595, 597, 618, 626
BERNSTEIN, F., 127
BEST, G. N., 83
Birds, 414, 533, 535, 536, 542, 545, 546, 548, 558, 560–565, 568
  Production of lens antibodies, 414
BLACKBURN, K. B., 78, 92
Blackhul kafir, 293, 297
BLAKESLEE, A. F., 87, 92, 93, 460, 463, 483, 485
BLANCHARD, F. C., 346
BLANK, E., 54
BLASIUS, 163
BLIZZARD, W. L., 548
Blood relationship of mice, 397
Blue green algae, 144
BOL, C. J. A. C., 548, 549
BONNIER, G., 212, 213
BORDET, J., 398, 406
BOSTIAN, C. H., 659
Botanical Garden at Amsterdam, Holland, 86
Botrytis, 194, 196
BOULENGER, G. A., 75, 83
BOYDEN, A. A., 415
Brassica, 450
  oleracea, 484
BREE, 50
BREWBAKER, H. E., 624
BRIDGES, C. B., 57, 105, 212, 284, 355, 375, 386, 389, 490, 497, 630, 633, 636
BRINK, R. A., 190, **595**, 596, 601, 614, 623, 624
BROOKE, H. C., 43, 50, 65
Brother-sister mating, 357, **360**, **364**, 371, 373, 374
Bruigera, 589
  Vivipary, 589
BRUNSON, A. M., 623, 625

Buchholz, T. J., 463
Bucknell University, 574
Burnham, C. R., 596, 597, 600, 610, 623–625
Burton, R. H., 506
Bush, B. F., 84
Bussey Institution, **42**, 43, 62
Buxton, B. H., 209, 450

Calder, A., 111
California Institute of Technology, **325**, **375**, **386**, 490, 505, **629**, 630
Call, L. E., 352, 356
Cameron, D. R., 196, 315, 485
Cammidge, P. J., 56
Campanula, 460
  Tetraploid, 460
Campbell, A., 50
Canary, 536, 537, 543
  Eye color
    Black, 536, 544
    Brown, 544
    Chocolate, 536
    Pink, 536, 544
  Linkage, 543
  Pigmentation, brown, 543
  Plumage color
    Chocolate brown, 536
    Cinnamon, 536, 544
Carnegie Institution of Washington, 225, 591
Carnoy's fluid, 597
Carotin, 299
Carpenter, D. C., 416
Carrier, L., 5
Carr-Saunders, A. M., 414
Cartledge, J. L., 87, 92, 93
Castle, W. E., 43, 45, 62, 72, 99
Cattle, Shorthorn breed, 111
Ceriops, 589
  Vivipary, 589
Chaerocampa elpenor, 484
Chalmers, R. E., 678
Chiarugi, A., 478
Chicken, 398, 402, 413, 414
Chipman, R. H., 192
Chlorophyll characters in sorghum, inheritance of, **291**, 293, 304–306
Christie, W., 536, 537, 539, 544, 548, 550, 552, 557–561
Christoff, M., 92
Chrom-acetic-formaldehyde solution, 424
Chromic acid, 645
Chromosome, split, **349**, 350, 352, 353
Citrus, 478
  aurantifolia, 478
  Triploid, 478
Class frequencies, equation of equilibrium of, 133
Clausen, J., 99, 450
Clausen, R. E., 92, 192, 209, 446, 449, 450
Cleistostoma ambigua, 588
Cleland, R. E., 325–327, 344, 345
Clements, F. E., 80, 93
Cobb, F., 345
Coccids, 251
Cocks, W. P., 56
Cole, L. J., 532, 536, 537, 542, 544, 548–552, 555, 556, 558, 559, 565
Collins, G. N., **4**, 604, 608, 625
Collins, J. L., **445**, 450
Color characters, 138
Color standards and nomenclature, 535, **544**, 545
Columba livia, (See pigeon) 551
Columbia University, 505
Conklin, 288
Conner, A. B., **291**, 292
Coolidge, 256
Coolidge tube, 661
Cooper, D. C., **595**, 597
Cope, 98
Copeland, 59–61, 72
Corn (See maize)
Cornell University, **175**, **490**, 599, 624, 625
Correns, C., 287, 288, 586
Cotton, 291, 300
Cousin matings, 373
Crane, M. B., 92–94
Crepidula, 288
Crépin, F., 86
Crepis, 92, 196, 443, 450, 484
  biennis, 445
  capillaris, 443, 484
  setosa, 445
  tectorum, 443
  Triploid, 484
Cresol, 401
Crew, 65
Crocker, W., 90
Crocker Institute of Cancer Research, 505
Crossbreeding, 146, 147
Crossing over, double, **372**, 373
Cryphaea grecillinia, 588
Cryphaea macrospora, 588
Cryptocoryne, 589
Cuénot, L., 43, 45, 47, 48, 54
Curves, distribution, 134
Cytology
  And genetics of crosses between fourteen- and seven-chromosome species of wheat, 309
  Identification of chromosome associated with the *v-g* linkage group in maize, 175
  Observations on Fragaria, 431
  Studies on Fragaria, **422**, **431**, 462

Daphnids, 251
Darlington, C. D., 92, 93, 342, 445, 446, 477, 479, 484
Darrow, G. M., 90, 437, 460
Darwin, C., 98, 99, 154, 537, 559
Datura, 87, 92, 93, 279, 460, 463, 483, 485, 597
  Diploids, 93
  Globe mutant, 483
  Tetraploid, 93, 460
  Triploid, 463
  Wrightii, 479

DAVENPORT, C. B., 591
DAUVART, A., 43, 52
DAVIS, B. M., 326–328, 346, 347
DAVIS, N. F., 574
DEMEREC, M., 284–286, 291, 295, 306, 602, 622
Dicnemon, moss, 588
DIGBY, L., 78
Digitalis, 450
  Hybrids, 209
Dihydroxyindole, 567
Di-hydroxyindole-2-carboxylic acid, 567
Dihydroxyphenylalanine, 567
Diptera, 252, 346
Distribution curves, 134
Distribution of gene frequencies and its consequences, 111, 133
DOBROVOLSKAIA-ZAVADSKAIA, N., 54
DOBZHANSKY, T., 212, 346, 375, 490, 491, 494, 495, **629,** 630, 632, 634, 636, 637, 644, 650, 652
Dog, 412, 413, 415
  Serum, 412, 413
Dominance ratio, **137,** 156
Dormancy, 574–576, 578–581, 583, 586–590
DORSEY, M. J., 76, 78, 93
DOWNS, 414
DRIVER, E. C., 274, 278
Drosera scheme, 192, 209, 210
Drosophila, 57, 61, 105, 143, 152, **212,** 213, 228, 251, 262, 263, 268, 287, 306, **346,** 349, 355, 374, 375, **386,** 477, 479, **490,** 493, 505, 512, 528–530, 633, 677
  Attached-X, **376, 377,** 380–385
  Attached-X females, 495, 496, 500–504
  Bar
    And infrabar, composite genes of, 274
    Deficiencies, 283
    Locus, facet numbers determined by genes, 277
    Locus, mutation at, 281
    Over infrabar, dominance of, 280
    Series, mutations and origins of the genes of, 268
  Chromosome
    Broken in translocations, **633,** 634, 651, 653
    Migration of the episome, 284
    Reduction, 630
  Conjugation, 630
  Constitution of primary exceptions, 393
  Crossing over, **493, 495,** 496, 498, 500–504, 629–631, 634–637, 652
    Frequency, 652–654
    In females carrying duplication, 495
  Crossovers, 393, 394
    Association of, **392**–394
  Cytological maps, 629, 630, 651–653
  Cytology, **644,** 650, 652, 654
    Of the translocations, 644
  Dominance of bar over infrabar, 280
  Episome on the chromosome, migration of the, 284, 285
Drosophila (Continued)
  Exceptions
    Constitution of primary, 393
    Genotypic constitution of, 389
    Phenotypic distribution of, **388,** 391
  Facet number, 279
    Determined by genes at the bar locus, 277
  Fertility, 493, 494
  Forked reversions, **380,** 381
  Frequencies of homozygosis of factors in attached-X females, 375
  Frequency of translocations, 490
  Gene
    Abnormal abdomen, 284
    Antlered wing, 280
    Apricot, 377, 379–385
    Aristaless, 491–496, 500–504, 630–632, 634, 635, 638–643, 654–658
    Bar eye, 267, **268–274–277–280, 281–283,** 284,286, 287, 378, 387, 395
    Bar and infrabar, composite, 274
    Bar series, 268
    Beadex, 378
    Black, 266, 491–497, 500–504, 630–635, 639–643, 653, 655–658
    Bobbed, 393, 494
    Bristle, 497, 636
    Cinnabar, 266, 497
    Crossveinless, 380, 387, 389, 395
    Curly, 491–493, 630–632, 637, 639–641, 643, 645
    Curved, 491–496, 500–504, 630–632, 634, 636, 639–643, 651, 653, 655–658
    Cut, 375, 376, 380, 387, 389, 395
    Double-bar eye (See ultrabar), 269, 272–274, 276, 279
    Dumpy, 491–496, 500–504, 630–635, 638–643, 651, 655–658
    Ebony, 490
    Echinus, 376, 377, 379–385, 387–389, 395
    Erect wing, 271
    Eyeless, 491, 630–632, 634, 635, 639–643, 655–658
    Forked, 271–273, 275, 281, 284, 375–378, 380–385, 387, 389, 393, 395, 493
    Full eye, 269–274, 276–279, 281–284, 287
    Fused, 271–273, 275
    Garnet, 375, 387, 389, 393, 395
    Hooked, 497, 498
    Infrabar eye, **273,** 274–276, **279, 280,** 281, 286
    Infrabar series, 279
      And facet number, 279
      Mutations and origins, 273
    Light, 497, 498
    Lobe, 637
    Mutable
      Magenta eye color, 285
      Miniature wing, 285
      Reddish-alpha, 285, 286
      Reddish body color, 285
    Mutability, 285
    Nick wing, 280

Drosophila (Continued)
Genes (Continued)
Notch, 262, 280, 284
Pink wing, 497
Plexus, 490–496, 500–504, 630–632, 634–636, 638–643, 651, 653, 655–658
Position effect, **275**–277
Purple, 266, 491–498, 500–504, 630–632, 634–636, 639–643, 651, 653, 655–658
Red eye, 271
Rolled wing, 497
Round eye (See full), 269, 273, 274
Ruby, 377, 379–385
Scraggly, 497
Scute, 254–256, 258–260, 262–264, 286, 376, 377, 379–385
Scute-1, 254–256, 258–266
Scute-2, 254–256, 258–266
Scute-3, 254–256, 258, 260–266
Scute-4, **256**–266
Scute-5, 255, 265
Scute-6, 255, 260, 262, 265
Scute-7, 263, 265
Sepia, 286
Sexual dimorphism, 275, **276**, 277
Speck, 490–496, 500–504, 630–636, 638–643, 651, 655–658
Strap wing, 280
Tan, 375, 376
Thick, 497
Ultrabar eye, 267, 269–274, 276, 277, 279, 281–283, 287
Vermilion, 380, 387, 389, 395
Vestigial wing, 280
White, 284
Wild-type, 268–271, 280, 284
Yellow, 255, 256, 262, 263, 285, 286, 376, 377, 379–385, 387
Genetical determination of size of duplicated piece, 497
Genic mutations, 505
Genotypic constitution of exceptions, 389
Gonads, **212–214, 215,** 218–**220**–224
Growth of, **212,** 213, **215–220**–224
Larval, 214
Heterosis, 378
Homozygosis, **375**–381, 388, 389, 391, 393–395
Inbreeding, 378
Infrabar, composite genes of bar and infrabar, 274
Infrabar, dominance of bar over, 280
Infrabar series, origin and mutation of genes, 273
Larvae, 213–220, 222, 223
Larval and pupal gonads, general characteristics of, 214
Linkage, 634
Of characters, 632
Of genes, 630, 631, 633, 653
Group, 651, 653
Loci at which the second chromosome is broken in translocations, **633,** 634, 651, 653
Drosophila (Continued)
melanogaster, **212,** 213, **254,** 257, 266, 268, 270, 279, 280, 285, 346, **375,** 381, 386, 490, **629,** 630, 645, 653
Migration of the episome on the chromosome, 284, 285
Mutable genes, **284**–286
Mutability of new formed genes, 285
Mutation, 505, 529, 530
And origin of the genes of the bar series, 268, 273
At the bar locus, 281
Somatic manifestation, 275
Non-disjunction, **638,** 641–644, 650
After X-ray treatment, 387
Second chromosone, **638,** 641, 643
Origin and mutation of the genes of the bar series, 268, 273
Origin of translocations, 630
Phenotypic distribution of exceptions, **388**, 391
Primary exceptions, **386**–389, 393
Primary exceptions obtained after X-ray treatment, 386
Protosome, the migration of the episome on the, 285
Pupae, 213–217, 220–222, 224
Roentgenization, 256, 262
Second chromosome, 493
Somatic manifestation of mutations, 275
Step allelomorphism in, **254,** 255, 262–265
Sterile, 494
Synthesis of attached-X line, 376
Transgene
Dachsous-1, 265
Dachsous-3, 265
Scute-4, 256
Characteristics and location of the gene, 256
Spoon, 262, 263, 265
Translocation, 346, **490–493**–496, 498, 500, 501, 503, 504, 512, **629, 630,** 631, 633, 634, 636, 648, 650–654
Cytological characteristics, 490
Cytology, 644
Effect of, on crossing over in the second chromosome, 493
Frequency of, 630
Genetical characteristics of, 490
Homozygous, **637,** 638, 640–642, 645, 648, 650, 653, 654
Involving the second and fourth chromosomes, **629**–632, 653
Loci at which the second chromosome is broken, **633,** 634, 651, 653
Origin of, 491
Pale, 490
Triploid intersexes, **212**
virilis, **284,** 285
Wild-type, 255, 256, 258, 260, 264, 265
X-ray radiation, 505, 528
X-ray treatment, **386, 387**–389, 393, 395

Drosophila (Continued)
X-ray treatment, frequency of primary non-disjunction after, 387
DUBININ, N. P., 254, 255, 262, 263
DUBOSCQ, O., 53
Duck, 398
DUNN, L. C., 44, 47, 48, 65, 72
DUNNING, W. F., 505
Duplication, complete, 144
DURHAM, F. M., 46–48, 536, 537, 544

EAST, E. M., 19, 60, 76, 94, 99, 145, 192, 422–424, 440, 441, 451, 453, 456, 457, 463, 465
Eel, 398
Eggplant, **1**, 2, 6, 9, 11, 13, 14, 17, 19, 20, 22–24
Branching habit, 19
Calyx spine, 20
Characters, **19**, 24
Color of fruit, 19
Crosses, 20
Increased yield of, **16**, 24
Relation between yield, **14**, 24
Relation of characters to increase in yield, 16
Cross-pollination, immediate effect of, **2**, 23
Cross-pollination, technique of, **22**, 24
Diameter of stem, **6**, 11, **13**, 23, 24
Diameter of stem and height of plant, relation between, **13**, 24
Effect of cross-pollination on the size of seed, 2
Expression of characters, **19**, 24
First-harvesting time and the yield, relation between, **19**, 24
Flowering habit, 22
Fruit
Color, 19
Shape, 19
Harvesting time, **13**, **19**, 24
Height of plant, **6**, 11, **13**, 23, 24
And diameter of stem, **6**, 11, **13**, 23, 24
Increase in, 23
Hybrid, **1**, 2
Vigor, **1**, 6, 19
And its practical utilization, 1
Increase in plant height, 23
Leaflet size, 19
Mechanism of flowers, 22
Parents, relation between yield, **14**, **16**, 24
Plant height, **6**, 8–11, **13**, 23, 24
Relation between the first-harvesting time and the yield, **19**, 24
Relation of characters to increase in yield, 16
Shape of fruit, 19
Size of leaflet, 19
Size of seed, effect of cross-pollination on the, 2
Stem diameter, **6**–9, 11, **13**, 23, 24
Technique of cross-pollination, **22**, 24
Variety, 3, 18
Kintyaku, 21
Sinkuro, 21
Sironasu, 21

Eggplant (Continued)
Yield of parents and that of crosses, relation between the, **14**, **16**, 24
Yield, relation between the first-harvesting time and the, **19**, 24
EIMER, 98
Eink rn, 309
Emergent evolution and creative, 154
EMERSON, R. A., 190, 291, 295, 605, 606, 608, 623–625
EMERSON, S., **325**, 327, 332
Emmer, 309, 312, 314–316, 322, 323
Environmental pressure, 146
Enzymes, Willstätter's structure of, 287
Equation of equilibrium of class frequencies, 133
Equilibrium, 100, **102**, 103, 105, 112, 116–118, 122, 126, 128, 129, 131, 133–135, 137, 138, 140–142, 146, 147, 149, 153, 155–157
Class frequencies, equation of, 133
Complete, 122
Frequencies, 104, 118
Simple Mendelian, 100
Under selection, 102
ERLANSON, C. O., 347
ERLANSON, E. W., **75**, 78–80, 86, 91, 92
Ethyl alcohol, 299
Eucalptodon, moss, 588
EVERSMANN, 160, 164
Evolution, **97**, 98, 100, 107, **142**, 143, 145–**147**, 149–**151**–**153**, **154**, 157, 158
Classification of the factors of, 142
Control of, 151
Creative and emergent, 154
In Mendelian populations, 97
Isolation, effect of, 98
Lability as the condition for, 147
Mean and variability of characters, 139
Of Mendelian systems, 142
Statistical factors
Fecundity, 98
Survival, 98
Theories of, 97
Evolutionary factors, 133
Evolutionary pressure, 105
Evolutionary process, 104, 105, **144**, **146**, 149, 150, 154
EYSTER, W. H., 288, **574**, 575, 584, 586, 612

Factor frequency, 103
Factors, evolutionary, 133
FARNHAM, M. E., 460
FEDERLEY, H., 99, 209, 484
Fegatella, 588
Fertility, effect of X-ray radiation on Habrobracon, 505
FINLAY, 402, 414
FISHER, R. A., 65, 69, 102, 106, 107, 109, 112, 115, 116, 119–121, 129, 132, **133**, 137, 138, 149, 156, 304, 567
FITZGERALD, 402
Fixation, 151, 156, 157
Complete, 117, 157

Fixation (Continued)
Of genes, 150
Random, 150
Rate of, 114, 115, 118
Flemming's fluid, 424, 645
Fly, 213, 225, 226, 228, 230, 236, 239, 240, 247–250, 270, 272, 281, 287, 490, 497, 630, 633, 634, 637–639, 642, 646, 650, 653, 654
FOCKE, W., 192
Fortunella hindsii, 478
Fortunella margarita, 478
FORTUYN, A. B. D., **160**, 161, **168**, **591**, 593
FOSTER, M. E., 349, 353
Fowl, 398
Fowl, domestic, 562, 564
Fragaria, 90, **422–431**, 441–445, 447, 449–451, **455**, 456, 460, 463, 465, 473, 485
Aberrant chromosome types, **465**, **467–469**, 470, 472, 474, 481–483, 486, 487
Descriptions, 465
Frequency and distribution, 470
Origin, 465
alba, 458
americana alba, 423, 424, 441, 450, 451, 457, 460, 461, 473, 474
Aneuploid forms, **455**, 464, 484
bracteata, 423–**426**, **428**, **430**, 431, 436–438, 441–443, 445, 451, 456, 457, 460, 461, 466, 468, 470, 471, 483, **485**, 487
californica, 423, **428**, 441, 451
Chesapeake, 444
chiloensis, 423, **424**, **428–430**, **431**, 434, 436, 438, 440, 442, **444**, 445, 447, 449–452, 461
Chromosomal chimera, 440, 456
collina, 423, **425**, **426**, **427**, 428, **430**, **431**, 434, 436, 441–445, 451, 461, 474
Crosses
Diploid and octoploid, **428**, 450–452
Hexaploid and octoploid species, 431
Octoploid and diploid, **430**, 444, 452
Octoploid species, 431
Seven-chromosome species crossed, 424
Seven-chromosome and twenty-one-chromosome species, 427
Cytological observations, 431
Diploid and octoploid crosses, 431
Octoploid and hexaploid crosses, 437
Somatic chromosome numbers, 437
Temperature data, **436**, 452
Cytological studies, **422**, **431**, 462
Description of crosses, 424
Diploid, 456, 460, 461, 463, 464, 466–468, 470, 473, 481–483, 485–487
Sterility of, 487
Docteur Nicaise, **444**
Dunlap, 428
Dwarfness, 469
Early Jersey Giant, 444
elatior, 423, 424, **427**, **428**, **431**, 437, 438, 442, 443, 452, **461–463**, 486
Faux hybrides, 422
Fertility, 456, 468, **480**
Fragaria (Continued)
Gene
Pink flower, 425, **448**, 457–459, 462, 466, 486
White flower, 458
Genetic studies, 422
glauca, 423, **429**, **431**, 432, 436, 438, 440, **448**, 449
grandiflora, 438, **445**
hagenbachiana, **445**
Hexaploid group, 456
Hunterdon, 462
Hybrid cytology of species crosses, 422
maxima, 423, **425**, **426–428**, **430**, 431, 434, 436, 441–443, **447**, 451
Meiosis, 476
Meiosis in triploid, 476
mexicana, 423, 441, 451
nilgerrensis, 423, **426**, **427**, **428**, **430**, 431, 437, 441–445, 451, **461**
Octoploid group, 456, 463
Out-crosses, 456, 462, 486
platypetala, 423, **424**, **431**
Polyploid forms, **455**, **464**
Species hybridization, 441
Sterility, 485, 487
Study of certain polyploid and aneuploid forms, 455
Tetraploid, 456, 459–**461**, 463–466, 468, 470–474, 476, 477, 483, 485, 486
Family, 456
Changes in compatibility with chromosome doubling, 463
Chromosome numbers, 463
Crosses between seven- and fourteen-chromosome groups, 460
Cytology of crosses, 462
elatior×tetraploid and reciprocal, 461
Inheritance in the tetraploid, 457
Octoploid and tetraploid crosses, 461
Origin, 456
Reduction division, 457
Triploid, 456, 465–468, 471–473, **476**, 477, **480**, 482, 483, 485–487
Derivatives, 480
Causes of variation, 485
Distribution of chromosomes, 483
Fertility in the triploid, 480
Progeny, 480
Segmental interchange between non-homologous chromosomes, 486
Substitution of chromosomes, 485
vesca, 422–**425**, **426–428**, **429**, **430**–432, 434, 436–438, 440–445, 447–452, 457, 460, 461, 465, 466, 468–**471**, **485**, 487
vesca rosea, 423, **425**, 438, 440, 441, 448, 451, 456
virginiana, 423, 424, **428**, **429**, **431**, 432, 434, 436–438, 440, 442, 444, 447, 449–452, 461, 464
Frequency
Equilibrium, **104**, **118**

Frequency (Continued)
Factor, 103
Random variation of gene, 106
FRIEDBERGER, 413
FRIEDLANDER, 414
FROST, H. B., 473
FULTON, R., 533, 537
Funaria, 374
Fungi, 588
Dormancy, 588

Gagea lutea, 87
GAHRINGER, J. E., 412, 413
GAINES, E. F., 446
GAIRDNER, A. E., 460
GAISSINOVICH, 255
Gall midge, 252
Galton and Mendel Memorial Fund, 225, 357
Gametes, random sampling, 111, 117, 119, 121, 126
GÄRTNER, C. F., 192
GASKOIN, J. S., 50
GATES, R. R., 478
GATES, W. H., 43, 47, 52–54, 59, 60, 68–70, 160, 166, 398, 400, 408, 592
GAY, 402
Gazzi magnani, 559
GEERTS, J. M., 478
Gene
At the bar locus, facet numbers determined in Drosophila, 277
Attraction and repulsion phenomena, 286
Autosomal, **364, 368,** 371, 373
Direction and frequency of mutations, 268
Dominance, 283
Episome, 268, 272–281, 283, **284,** 285–288
Frequency, **100–106,** 107, 110–114, 121–123, 125–131, 134–139, 141, 142, 148–151, 155–157
Distribution of, and its consequences, **111,** 133
Variation, 100, 106
Intermediate variegation, 288
Linked, 374
Loss of, 114
Mutation, 99, 142–145
Structural explanations of, 287
Origins and mutations of the bar series in Drosophila, 268, 273
Protosome, 268, 272, 274–280, 283–**285,** 286
Quantitative series with a completely dominant allelomorph, 279
Random fixation, 150
Repulsion phenomena, attraction and, 286
Sex linked, **360, 367,** 371, 373, 374
Side chain, **267,** 268, 273, 275, 276, 284–286
Specificity, 142
Structure, **267,** 272, 275, 276, 281, 283, 287, 288
Symmetry, 286
Unlinked, 367, 368
Genetic studies on Fragaria, 422
Genetics and cytology of crosses between fourteen- and seven-chromosome species of wheat, 309
GENGOU, 398
GHIGI, A., 559, 560
GILMORE, K. A., 659
Goat, 398
GOEBEL, K., 589
GOLDSCHMIDT, R., 251
Gonads in Drosophila, **212,** 213, **215–220–**224
GOODSPEED, T. H., 92, 94, 99, 192, 450, 479
GORDON, G., 52
GOWEN, J. W., 376, 479, 505, 529
GOWEN, M. S., 376
Gramineae, 306
Grapes, 76
Grassh ppers, 353
GREGORY, R. P., 460
GRIFFEE, F., 5
GRIFFITHS, E., 347
Growth of the gonads in Drosophila, **212,** 213, **215–220–**224
Guinea pig, 138, 279, 398, **400,** 401, 407, 410, 413, 415
GUYER, 414

Habrobracon, **505,** 506, 529, **659,** 676, 677
Abnormal fertilization, 513
Abnormal individuals, **505, 507,** 509, 512
Abnormalities
Inherited, 675
Not inherited, 666, 674, 677, 679
Broad abdomen, 666
Notched wings, 666
Binuclearity of the egg, 660, 678, 679
Character
Abnormal abdomen, 664, 669, 674, 677
Abnormal abdominal sclerites, 664, 666
Abnormal genitalia, 664, 674
Abnormal venation, 674
Antennae abnormal, 664, 674, 677
Antennae bent, 674
Antennae deformed, 664
Antennae twisted, 674
Antennal segments abnormal, 664
Antennal segments fused, 664, 674
Beaded, 665, 667
Black eye, 677
Black-ivory eye, 677
Black and orange eye, 668, 675, 677
Cantaloup, 677
Clubbed antennae, 674
Confluent, 675
Disorganized veins, 668
Double right primary wing, 665
Extra wing, 674
Feet abnormal, 664
Ivory and black eye, 666
Ivory ocelli, 666
Kidney eyes, 676
Legs deformed, 664
Median ocellus small, 664, 677
Mesosternum, 662

Habrobracon (Continued)
Character (Continued)
Ocelli, 674
Ocelli light, 668
Primary wing short, 669
Primary wing small, 674
Pronotum, 674, 675
Reduced wing, 669
Short wing, 675
Shot, 677, 679
Shot wings, 671–673
Small eyes, 665, 674
Sting abnormal, 664, 677
Stumpy, 675
Truncate, 675
Tumor on eye, 669
Wing veins disorganized, 664, 671, 672, 679
Wings wrinkled, 664, 669, 674
Yellow basal antennal segments, 665
Defective individuals, 668, 677
All ocelli absent, 668
Double foot, 668
One antenna absent, 668, 677
One eye absent, 668, 669, 674
Effect of X-radiation of larvae, 659
Effect of X-ray on viability of the treated individuals and their descendants, 527
Effect of X-ray radiation on fertility, 526
Effect of X-ray radiation on occurrence of abnormal individuals, 505, 507
Effect of X-ray radiation on occurrence of fertility, 505, 526
Effect of X-ray radiation on occurrence of mutation rate, 505, 507
Effect of X-ray radiation on occurrence of viability, **505, 527,** 528
Fertility, **505,** 507–509, 512, 516–520, 523, 524, **526**–530
Freaks, 664, 666, 676, 679
Gene
Antennae fused, 511, 514, 525, 530
Bar eye, 506, 512–514, 518–525, 529, 530, 674, 675
Black eye, 506, 511–514, 516, 517, 520, 524, 529
Cantaloup, 677
Cream-colored eyes, 525
Defective median vein, 511, 514, 525, 530
Defective wing venation, 520, 521, 524, 525, 530, 531
Eye-color, 506, 512
Eyeless, 506, 511, 512, 514, 519, 521–525, 529, 530
Ivory eye, 506, 512–514, 516, 517, 529–531, 660–662, 664, 666, 668, 669, 675–678
Ivory eyes and ocelli, 280
Light ocelli, 280
Normal eye, 522, 524
Notched wings, 525
Orange eye, 506, 509, 511–514, 516–518, 520, 524, 529–531, 660, 662, 666, 668, 675–678
Habrobracon (Continued)
Gene (Continued)
Orange eyes and ocelli, 280
Red eye, 525
Reduced wing, 506, 509, 512, 513, 516, 518, 520–524, 529–531, 660–662, 664, 666, 668, 671–673, 675–679
Shot, 512
Small, 523
Vestigial wings, 511, 514, 525, 530
Wild-type, 280
Wing, normal, 506, 513, 514, 516, 518, 520, 523, 524
Genetic changes, 530
Gynandromorph, 678
Iowa City stock, 506
juglandis, 280, **505,** 506, 530, 659
Mosaic, 660, 665–669, 671, 673, 675–679
Mutant, 675, 676
Beaded, 665, 667
Confluent, 675
Ivory eye-color stock, 506
Kidney eyes, 676
Miniature, 669
Orange eye-color stock, 506
Reduced wing stock, 506
Stumpy, 675
Truncate, 675
Mutation, 660, 665, 667, 669, 673, 675–679
Antennal segments, 665
Rate, **505–507**–509, 512, 513, 517–520, 523–525, 529, 530, 678
One-eyed forms, 677
Somatic abnormalities, 507, 508, 510, 511, 526, 529, 530
Sterility, 508, 516–518, 520, 523–525, 527
Translocations, 512
Viability, **505, 527,** 528, 530
X-ray radiation, **505–507,** 509–512, 516–518, 520, 523–**526, 527**–531, **659**
HADLEY, P. B., 405, 420
Haemotoxylin, 208
HAGEDOORN, A. C., 107
HAGEDOORN, A. L., 107
HAGIWARA, T., 32, 36
Haidenhain's iron-alum-haematoxylin, 424
HAKANSSON, A., 319, 434, 626
HALDANE, J. B. S., 102, 142, 149, **357**
HALTER, C. R., 55
HAMBURGER, F., 398
HANCE, R. T., 45
HANSON, F. B., 85, 270, 273, 529
HARDY, G. H., 100
HARRINGTON, J. B., 300
HARRISON, J. W. H., 78, 92
HARRISON, R. G., 544
Harvard University, 42, 422, 455
HAWKINS, L. E., 532, 542, 544, **547,** 558
HAYES, H. K., 14, 17
HEILBORN, O., 87
Hematoma, 402
Hereditary variation, 98
Heredity, 97, 98, **144,** 153, 154

HERIBERT-NILSSON, 339
HERSH, R. K., 274, 276–278, 287
Heterozygosis, **107**–112, 115–117, 139, 141
  Loss of, **107**, 109–112, 116, 139, 141
  Rate of decrease, **107**, 109, 110, 112, 116, 139, 141
HICKS, R. A., 397
HILL, H. E., 175
HOLLINGSHEAD, L., 312, 443, 445, 450
HOPPE, E. N., 227
Horse, Clydesdale, 111
HOWARD, H. A., 56
HUETTNER, A. F., 213
HUNT, H. R., 52, 54
HURST, C. C., 75, 82–86, 94, 449
HUSKINS, C. L., 75, 92, 479
HUTCHISON, 306
HUXLEY, J. S., 414
Hyacinths, 484
  Triploid, 484
Hybrid, 99, **191**
  Digitalis, 209
  Interspecific, 191
    Chromosome behavior, 191
    Chromosome constitution of F_2 progeny, 191
    Chromosome constitution of functional male and female gametes, 191
    Crossing over, 191
    Meiotic behavior of chromosomes, 191
    Recombination possibilities, 191
  Pygaera, 209
  Raphanus-Brassica, 208, 209
  Species, 480
  Sterility in roses, 75
Hybrid vigor, 1, 6, 17, 19, 373
Hybrid vigor in eggplants and its practical utilization, 1
Hybridization, 99
  External, 146, 147
  Interspecific, **191**, 192, 210
Hymenoptera, 671

ICHIJIMA, K., 441, 456, 457
IDE, S. I., 401
ILLICK, J. T., 326
Illinois Natural History Survey, **267**, 289
IMAI, Y., **26**, 29, 30, 33, 36, 39, 47
IMMER, F. R., 304
Immigration, 148
Inbreeding, 106, 107, 110, 111, 141, 146, 277, 292, **357**, 367, **370**, 372–374
  And linkage, 357
  With initial population, 370
Inheritance, 98, 102
  In house mouse, 42
  In pigeons, 532, 547
  Mode of, 102
  Of acquired characters, 98
  Of certain characters in Oenothera hybrids of different chromosome configurations, 325
Inheritance (Continued)
  Of chlorophyll characters in sorghum, **291**, 293, 304–306
  Of vivipary in maize, 578
Intensity of selection, 135, 136
Interspecific hybrid
  Chromosome behavior, 191
  Chromosome constitution, 191
  Chromosome constitution of functional male and female gametes, 191
  Crossing over, 191
  Meiotic behavior of chromosomes, 191
  Recombination possibilities, 191
Interspecific hybridization, **191**, 192, 210
Interspecific sterility, 191, 193, 195
Iodine-crystal violet, 424
Iowa State College, 624
Iron haematoxylin, 193, 645
Irreversible recurrent mutation, 128
Isolated population, 114, 116
Isolation, 98, 99, 114, 116, 137, 150, 153, 158

Japanese waltzing mouse, 398, **400**, 408, 411, 413, 417, 419, 420
JAYNE, H., 69
JEFFREY, E. C., 75
JENKINS, J. A., 446
JENKINS, M. T., 190, 610, 612, 625
JENNINGS, H. S., 106, 109, 357, 358, 363, 367, 368
JOHANNSEN, 99
Johns Innes Horticultural Institution, 357
JONES, D. F., 5, 14, 17, 19, 102, 616
JONES, S. v. H., 548
JORGENSON, C. A., 93, 446
Journal of Experimental Biology, 254

Kafir (See sorghum), 291–294, 297, 299, 301, 302, 307
Kahn test, 406
KAKIZAKI, Y., 1
KANO, T., 39
Kansas Agricultural Experiment Station, 349, 353
Kansas State Agricultural College, 349, 353
KARPECHENKO, G. D., 208, 209, 450, 484, 644
KARPER, R. E., **291**, 292
KEELER, C. E., 48, 52, 53, 56, 72
KELLEY, F. J., 532, 542, 544, 548–552, 555, 556, 558, 559, 565
KEMP, W. B., 152
KEMPTON, J. H., 4, 608, 624
KENDALL, J., 87, 92, 480
KERKIS, J., 212
KIDA, M., 9, 13, 14, 17
KIHARA, H., 310, 446
Kihara's scheme, 198
KIRKHAM, W. B., 59
KÖLREUTER, 192
KOSTOFF, D., 87, 92, 479, 480
KRAFKA, J., 274, 276, 278
KRISTOFFERSON, K. B., 86, 89

KULKARNI, C. G., 326, 346
KVAKAN, P., 610

Lability as the condition for evolution, 147
LADEBECK, E., 564
LAMARCK, 98, 142
Lamarckism, 99
LAMMERTS, W. E., **191**, 196, 208, 450, 484
LANDAUER, W., 544
LANG, A., 53
Leaf mold, 228
LEBEDINSKY, N. G., 43, 52, 62, 72
Lebistes, 374
Leitz apochromatic oil immersion lens, 193
Leningrad Academy of Science, 212
LESLIE, J. W., 92, 484
LI, 610
LINDSTROM, E. W., 291, 295, 300, 575, 582, 624
Linkage, **357**, 363
  Groups, association of semisterile-1 in maize, 595
  Groups, in maize, **175**, 178–180, **186**, 187, 189, 190
  Of swollen and narrow in Sciara, 247
  Studies in Pharbitis, 26
Linked genes, 374
LITTLE, C. C., 44, 46, 54–57, 68, **397**, 399
Liverworts, 588
  Germination, 588
  Vivipary, 588
LLOYD-JONES, O., 552, 562–564
LONGLEY, A. E., 78, 93, 440, 460, 478
LORD, E. M., 53
Loss of genes, 114
LOTKA, A. J., 98
LOTSY, 99
LUCE, W. M., 274
Luer, tuberculin, 399
LUMLEY, W. F., 537
Lycium halimifolium, 87, 479, 480
Lycopersicum, 92
LYELL, J. C., 533–535

MCCLINTOCK, B., 92, **175**, 177, 437, 473, 476, **484**, 622
MACDOWELL, E. C., 591, 593
MCNABB, J. W., 353
MCPHEE, H. C., 107, 111
MAHONEY, C. H., 300
Maize, 4, 5, 17, 288, 291–293, 295, 300, 305–307, 529, 574–**575**–**578**, 584, 586–590, **595**–597, 624–626
  Aborted pollen, 613, **614**–616, 622, 626
  Aborted pollen grains in semisterile plants, 614
  Albinism, 575
  Albinotic viviparous seedlings, 575
  Association of semisterile with linkage groups, 595
  Chlorophyll mutations, 306, 307
  Chloroplastid pigments, 575, 576
  Chromosome, distribution, 618
Maize (Continued)
  Chromosome ring, **597**–599, **618**–622, 625–627
  Crosses, increased yield of, relation between and the yield of parents, 17, 24
  Crossing over, **622**, 623
    Effect of semisterility on, 622
  Cross-pollination on the size of seed, effect of, 4
  Distribution of the chromosomes at the reduction divison, 618
  Dormancy, 575, 576, 578–581, 583, 586, 587
  Effect of cross-pollination on the size of seed, 4
  Endosperm defects, 575
  Fertility, 600, 603, 605, 606, 608, 609, 612, 618, 624
  Gametes produced by semisterile plants, 616
  Gene
    Adherent, 600, 608, 609, 618, 626
    Aleurone color, 579–**581**–**584**, 585, 599–**601** 602, **611**, 613
      Brown, 600, 609, 610, 613
      Purple, 580, **583**, **585**, **610**
      Red, **583**, 600, 610, 613
    Anther ear, 600, 608, 609, 626
    Brachytic, 600, **605**–607, 609, 618, 623, 626, 627
    Brevis, 600, 610, 613
    Brittle, 600, 610, 613
    Brown midrib, 600, 612
    Cream, 612
    Crinkly, 600, 611, 613
    Dormant embryo, 586, 587
    Dwarf, 599, 600, 610, 611, 613
    Endosperm
      Reduced, 580, **584**, **585**, **586**, 589, 590
      Scarred, 580, **584**, **585**, 590
      Shrunken, 579, 580, **583**, **584**, **585**, **586**, 587, 600, **601**, 613
      Sugary, 580, **583**, 586, 587, 600, **602**, 613
      White, 584
      Yellow, 580, **584**–**586**, 590
    Fine-striped, 600, 606–609, 623, 626, 627
    Glossy, **609**, 610, 613
    Golden, 580, 600, **601**, 602, **612**, 613
    Green, 603, 607, 608
    Green midrib, 612
    Iojap, 600, 609, 613
    Japonica, 600, **613**
    Liguleless, 600, **602**–604, 618, 623, 625, 627
    Orange pericarp, 288
    Pale green, 600, 611, 613
    Pericarp color, 600, **605**, 606, 618, 626
    Plant color, 580, 600, 605, **611**, 613
    Purple, 605
    Ragged, 599, 600, 601, 611, 613
    Ramosa, 600, **609**, 610, 613
    Silkless, 600, 602, 604, 605, 625
    Summary, 580
    Tall, 607, 609, 611
    Tassel seed, 599, 600, 605, 606, 608, **611**, 613, 625, 626

Maize (Continued)
Gene (Continued)
Tunicate, 600, **602**, 613
Virescent, 600–604, **610**, 613, 618, 623, 625, 627
Viviparous embryo, 577–581, 583, 586, 587
Vivipary-1, 578–**581**, **582**, **583**, **586**, 589
Vivipary-2, **578**, **583**, **584**, **586**, 589
Vivipary-3, **578**, **586**, 589
Vivipary-4, **579**, **587**, 589
Waxy, 600, 601, 613
Germinating seeds, 575
Germination, 575, 576, 578–580, 587
Golden Glow, 600
Hybrid vigor, 17
Inheritance of vivipary in, 578
Linkage, 575, 579, **581**–**583**–**586**, **587**, 589, **595**, 597, **599**–601, 609–**612**, 623, 625–627
Experiments, 599
False, 626
Groups, **595**, 600, 601, 609–612
Relations of vivipary, 581, 583, 586, 587
Tests, 627
With aleurone and plant color-tassel seed group, 611
With aleurone color-golden group, 601
With genes in the aleurone color-shrunken group, 601
With genes whose linkage relations are unknown, 612
With liguleless group, 602
With pericarp color-brachytic group, 605
With purple-aleurone-virescent group, 610
With ramosa-glossy group, 609
With sugary-tunicate group, 602
With yellow endosperm-purple plant group, 605
Mutations, chlorophyll, 306
Origin and description of vivipary in maize, 575
Physiological correlation, 575
Premature germination, 574–576, 578–580, 587, 588
Ratios of normal versus viviparous kernels, **579**, 580
Seedlings, green viviparous, 575
Semisterile, **595**–**597**, 598, 600, 605, 608, **614**, 615–620, 622, **623**, 625–627
Normal, 596–598, 605, 617, **623**–627
Normal class of plants from, 623
Occurrence of a chromosome ring in, **597**, 619, 620, 625
Plants, 614, 616
Semisterility, 595–597, 599–607, 609–613, 616, **622**, 625–627
Effect on crossing over, 622
Size of seed, effect of cross-pollination on the, 4
Spurious ratios of normal versus viviparous kernels, **579**, 580
Sterility, 596, 599, 617, 618, 625, 628

Maize (Continued)
Sun red, 605
Translocation, 596, 597, 608, 609, 617, 623, 626
Virescent seedlings, 295
Viviparous, 575, **579**–584, 586, 587, 589, 590
Embryos, increased moisture, 580
Embry s, premature drying, 579, 580
Kernels, spurious ratios of normal versus, **579**, 580
Vivipary, **574**, **575**–**578**–**581**–584, 586, **587**, 589, 590
Inheritance of, 578
Linkage relations, **581**, **583**, **586**, 587
Yellow seedlings, 301
Yield of parents and the increased yield of crosses, relation between the, 17, 24
Malva
crispa, 90
Hybrids, 86
Interspecific, 89
neglecta, 90
pusilla, 90
Mammals, 414, 532, 543, 548, 565, 568
MANGELSDORF, A. J., 94, 422–424, 431, 440, 441, 451, 456, 457, 463, 465
MANGELSDORF, P. C., 306, 456, 575, 578, 587
MANN, M. C., 445
MANWARING, W. H., 415, 416
Marine Biological Laboratory, 659
MARRYAT, D. C. E., 536, 537, 544
MARSHAK, A., 456
MARSHALL, C. C., 50
Mating
Assortative, 146
Random, 111, 112
System, 102
Matthiola, 473
MAVOR, J. W., 386
MAY, H. G., 268, 287
Meiosis, 145
Melanin, 567
Mendelian
Basis, 153
Equilibrium, 100
Heredity, 98, 99
Mechanism, 97
Populations, evolution in, 97
Recombination, 145
System, 106
Evolution of, 142
Mentzelia multiflora, 93
Mentzelia nuda, 93
Metopsilus porcellus, 484
METZ, C. W., 225–228, 230, 231, 234, 238, 241, 251, 252
METZELAAR, J., 398, 533–535, **537**, 541, 544, 547, 550–552
MEURMAN, O., 93
Michigan Academy of Science, Arts, and Letters, 387
Michigan Botanical Garden, 80

Michigan State Department of Conservation, 544
Migration, **100**, 106, **111**, 117, **126**, 135–137, 155
  Pressure, **100**, 134
  Rate, 135
Miles, F. C., 295
Millardet, M. A., 422, 441, 444, 449–451
Miner, J. R., 508, 511
Mitosis, 144
Miyazawa, 36
Mohr, O. L., 284
Morgan, L. V., 375, 376, 379, 386
Morgan, T. H., 44, 56, 57, 99, 105, 270, 273, 280, 284, 380, 505, 560, 630, 636
Morning glory, Japanese (See Pharbitis), 26
Moscow Zootechnical Institute, 254
Moses, M. S., 225, 227, 241
Mosses, 588
  Germination, 588
  Vivipary, 588
Mouse, **42–44**, 46, 47, 50, 52–54, 56, 57, 59, 63, 71, 72, 160–162, **397–400**, 401, 408–413, 415, 416, 418, 420, **591**, 593, 594
  Antigen, 398–**400**, **401**, 402, **405–407–419**
    Anti-Japanese waltzer rabbit sera, 417, 418
    Anti-mouse rabbit sera, 417, 419
    Anti-mouse sera, 418
    Anti-rat rabbit serum, 417, 418
    Chicken serum, 413
    Dilutions, **405**–408, 416, 418, 419
    Extract of guinea pig kidney, **400**, 401, 407, 410
    Extract of muscle tissue, **400**, 407–410
    Horse serum, 408, 411
    Human serum, 409–411, 413
      Caucasian, 408, 410
      Ethiopian, 408, 410
      Five different races, **401**, 408, 410
      Indian, 408, 410
      Malay, 408, 410
      Mongolian, 408, 410
    Mouse serum, 410, 411, 414, 418
    Mus musculus
      blood, **401**, 407, 409
      organ transfer, 408
      serum, 407
    Rabbit serum, **401**, 408–411, 413, 420
    Rat serum, **401**, 408–411
    Rodent serum, 411
  Antiserum
    In pigeons, 397
      Antigen index, 407
      Asepsis, 407
      Collection, **403**, 406
      Correlation, 414
      Precipitin test, 399, **405**, 407, 417
      Precipitin test, preparation of glassware, 405
      Preparation of antigen, 399
    In rabbits, 416
    Precipitating, **397**, 403, 406, 408–410, **416**
  Black, 591, 593
  Black-and-tan, probable mutation, **67**, 68
Mouse (Continued)
  Blood relationship, 397
  Body size, 165
  Body weight, 167, **168**–174
  Characters, 43, 56
  Correlation
    Between tailrings and body size, 165
    Between tailrings and tail length, 165
    Between tailrings and tail vertebrae, 166
  Dark-eyed intense black, 593
  Dominance of dilution, reversal of, 61
  European house, 160
  Gene, 58, 64
    Agouti, **43**–46, 57, 63, 68, 71
    Agouti and its allelomorphs, 43
    Agouti, white bellied, **44**, 68
    Albinism, **45**, 46, 53, 56, 57, 59, 63, 67, 71
    Angora, 56
    Black, 46, 48
    Black-and-tan, **43**–45, 57, **67**, 68, 72
    Black-eyed white, **47**, 53, 57, 59, 63, 71
    Brown, 45, **46**, 48, 59, 63, 71, 399
    Chinchilla, **45**, 57
    Chocolate, 62
    Cinnamon, 62
    Color, 45
    Congenital eye anomaly, 56
    Corrugated hairless, 50
    Density, 59, 61, 62
    Dilute, 62, 399
    Dilution, 43, **46**, 47, 52, 59, 61, 63, 71, 72
      Extreme, **45**, 57
      Linkage of short-ear and, 59
      Reversal of dominance, 61
    Dwarf, 55
    Ectromelus, 54
    Flexed-tail, 52, **54**
    Haemoragic head, **54**, 55
    Haired, 66
    Hairless, **42**, 43, 47–52, 59, **63**, 65, 66, 71
    Hyperglycaemia, **56**, 57
    Intense, 61, 62, 72
    Kinky-tail, 52, 68, 72
    Light-belly, 57
    Long-eared, 62
    Naked, **42**, 43, 49, **50**–52, 57, 59, **62**, 63, 67, 71, 72
      Linkage of, 42, 62
    Normal-ear, 59, 61, 69, 70
    Piebald, 43, **47**, 48, 50, 59, **63**, 65, 66, 71
    Pink-eye, 45, **46**, 53, 63, 67, 71
    Polydactylism, 54
    Rodless, 48, **52**, 53, 57, 63, 71, 72
    Sex-linked lethals, 56
    Shaker, 45, 46, **53**, 59
    Short-ear, **42**, 43, 47, **52**, **59**, 61–63, **68**–72
    Silver, 46, 48, 53, 57
    Tailless, **53**, 57, 59
    Twisted nose, 56
    Waltzing, **53**, 63, 71
    Yellow, **44**–46, 53, 57, 59
  Inbreeding, 170
  Inheritance, 42

Mouse (Continued)
Japanese waltzing, 160, 398, **400**, 408, 411, 413, 417, 419, 420, 592
Linkage, 42
Mendelian segregation, 594
Mutations to black-and-tan, 67
Pink-eyed dilute brown, 591–593
Precipitating antisera in pigeons, 397
Controls, 407
Observation and reading of tests, 406
Production of antisera, 402
Technique of precipitin test, 405
Precipitating antisera in rabbits, 416
Protein, anaphylactic shock in pigeons sensitized to, 412
Reversal of dominance, 61
Short-ear, effects of the gene causing, 68
Spotting modifiers, 47
Strain, StoLi, 591–593
Tail length, 165
Tail vertebrae, number of, 166
Tailrings, cross between, with different numbers of, 591
Tailrings, number of, **160**–170, 173, 174, **591**–594
Unit characters, 43, 56
Waltzing, 160, 166
MULLER, H. J., 57, 99, 143, 144, 213, 254, 255, 279, 349, 380, 460, 490, 505, 512, 528, 530, 623, 630, 631, 634, 637, 644
MURRAY, W. S., 399
Mus
bactrianus, 160, 398, 400, 401, 407, 409, 417, 419, 420
faeroensis, **400**, 407, 410, 417, 419, 420
gansuensis, 160
musculus, 53, **160**–**168**–174, 399, **400**, 401, 407–413, 417–420, 591, 592, 594
norvegicus, 408, 417–419
wagneri, 53, **160**–**168**–171, 174
wagneri manchu, 160
wagneri mongolium, 160
Mutable genes of Drosophila, 284–286
Mutation, 98–**100**, 103–107, **111**, 114, 116–**121**–126, **128**–138, 140–145, 148–153, 155–157
At the bar locus, 281
Direction and frequency, 268
Irreversible, 128–133
Migration or selection, 111
Nonrecurrent, 116
Pressure, **100**, 103, 104, 128, 134, 137, 139, 141, 142, 149–151, 156
Random, 142
Rate, 134, 135, 144, 157
Effect of X-ray radiation on Habrobracon, 505, 507
Recessive, 137
Recurrent, **121**, **128**, 155
Recurrent irreversible, 128
Reverse, **121**, 122, 129, 136, 137
Reversible, 132, 133
Reversible recurrent, 121
Somatic manifestation, 275
Mutation (Continued)
Structural explanation of gene, 287

NABOURS, R. K., 349, 352–356
NACHTSHEIM, H., 63, 72
NAGAI, K., 9, 13, 14, 17
NAGAO, M., 39
NÄGELI, C., 53, 98
National Research Council, 505, 595, 659, 661
Committee for Research in Problems of Sex 225
National Research Council of Canada, 323
NAVASHIN, M., 92, 484
Navashin's
Chrom-acetoformaldehyde solution, 477
Fluid, 645
Modified solution, 193
Nematodes, 251
NEWTON, W. C. F., 78, 209, 445, 446, 450, 473, 477, 484
NICOLAS, J. H., 78
Nicotiana, 76, 92, **191**, 192, 449, 450
Aceto-carmine mount, 201
Amphidiploid, **191**, 199–**204**–210
Crossing amphidiploids with parental species, 204
Derivatives obtained by selfing, 198
digluta, 209
Hybrid, amphidiploid, **191**, 199–**204**–210
Interspecific hybrid
Amphidiploid, **191**, 198–**204**–210
Amphidiploid lines, origin and behavior of, 198
Backcrossing, 191–198, **204**, 205, 208, 210
Chromosome behavior, 201, 202
Chromosome determination, 210
Chromosome number, 198, 199, 210
Derivatives, **198**, 201, 210
Dwarf types, 200, 202, 203
Fertility, 198, 199
Flower shape, 199, 205
Flower size, 199, 205
Giant types, 200, 206, 207
Infertility, 199
Morphology, 198, 199, 205
paniculata-rustica, 192, **193**–198, 200, 209, 210
rustica-paniculata, 195, **196**–198
rustica pumila, 198, 204
Sterility, 202, 203, 208, 210
Interspecific hybridization in, **191**, 192
Langsdorffii, 479
paniculata, 89, **191**–200, 208, 210, 484
Reciprocal backcrosses, 208
paniculata-rustica, functional gametes, 192, **193**
rustica, **191**–200, 210, 484
rustica pumila, 193–200, **204**, 205, 210
rusticana, 89
rustica-paniculata, **191**, 197
Sterility, 484
sylvestris, 450

Nicotiana (Continued)
sylvestris-Tabacum, 209
Tabacum, 450
Tabacum macrophylla, 479
NILSSON-EHLE, 99, 300
NISHIYAMA, I., 310, 478
NOLF, P., 398
North Carolina State College of Agriculture, 659
North Central College, 289
NOVY, F. G., 416
NUTTALL, G. H. F., 397, 398, 404, 415

Oats, 479
OEHLKERS, F., 325, 345
Oenothera, 102, **325**, 342, 344, 346, 478
Belling hypothesis, 342
biennis, 326, 345
Chromosome complex, 345
biennis sulfurea, 345
character, 325
Dwarf, 338, 340, 341, 347
Green bud-cones, 326, 332, 334, 336, 339, 340, 342, 347
Growth-habit, 326, 327
Red bud-cones, 326–332, 334, 336, 338–340, 344, 347
Revolute leaves, 345, 346
Splashed bud-cones, **327**, 338–340, 347
Stature, 330, 331, 334, 336, 338, 340, 341, 347
Sulfur-colored flowers, 326, 330, 331, 334, 336, 338, 340, 341, 345, 347
Yellow flowers, 326, 331, 336, 338, 340, 341
Chromosome complexes, 339
Chromosome configurations, 325, 332
Crosses between *franciscana sulfurea* and *franciscana*, 328
Crosses between *franciscana sulfurea* and sulfur-dwarf, 327
formosa, 345, 346
franciscana, **326–328**, 338–342, 344, 345, 347
franciscana sulfurea, **326**–328, 338–342, 347
Sulfur-dwarf, **327**–329, 334, 336, 338–340, 342, 344, 345, 347
Gametophyte competition, 341
gaudens, 345
genes, linkage of, 341
gigas, 478
Hookeri, 327
Hybrids, **325, 332**, 341, 347, 478
Inbreeding, 340
Lamarckiana, 345, 478
lata, 478
Linkage relationships, 325, **340**, 342
nanella, 345
Pollen grains
Empty, 326, 327
Shrivelled, 326, 327
Pollen lethal, 340
pratincola, 345, 346
rubens, 345
rubricalyx, 478
Oenothera (Continued)
Translocations, 342
Triploids, 478
velans, 345
OESTERLING, H. C., 289
Oklahoma Agricultural and Mechanical College, **547**, 548
Ontogeny, 288

PACKARD, C., 506
PAINTER, T. S., 53, 490
Paratettix, 350, **353**, 354
Parattix texanus, **349**–353
A split in chromosomes about to enter the spermatid, **349**, 350, 352, 353
Parent and offspring mating, **367, 368**, 371, 373, 374
PARKER, M. C., 624
PATTERSON, J. T., 529, 677
PAYNE, 254
PEARL, R., 508, 511
PEARSON, E. S., 56
PEARSON, K., 115, 119, 122
Peiping Union Medical College, 591
Peking Union Medical College, **160**, 161, 166, **168**
PELLEW, C., 78, 209, 473
Pellia, 588
Vivipary, 588
PENLAND, C. W., 80, 93
Pennsylvania College for Women, 659
PERCIVAL, J., 446
Petrol-ether, 299
Pharbitis Nil, **26**, 39, 40
Gene
Acuminate, 32, 40
Blown-1, 26, 40
Blown-2, 35, 41
Brown, 26, 27, 40
Buff, 38
Contorted, 39
Contracted, 32, 34, 40
Cordate, 28–30, 40
Couple, 28, 40
Cream, 32, 33, 40
Crêpe, 29, 30, 40
Crisscrossed, 38
Crumpled-1, 26–28, 40
Crumpled-2, 35, 41
Deficient, 38
Deformed, 27, 30, 31, 40
Delicate, 35, 41
Dilute, 32, 36, 41
Dragonfly, 32, 36, 37, 41
Duplicated, 36, 37, 41
Duskish, 32, 34, 40, 41
Dusky, 30, 31, 40
Even, 27, 28, 40
Extended, 37, 41
Faded, 28, 40
Fasciated-1, 35, 41
Fasciated-2, 35, 41
Fasciated-3, 26, 40

Pharbitis Nil (Continued)
Gene (Continued)
Feathered, 27, 28, 30, 40
Flecked, 39
Foliate, 38, 41
Glabrous, 38, 39
Intense, 32, 34, 40
Interaxil-green, 32–34, 40
Light, 30, 31, 40
Lined, 39
Magenta, 32, 40
Maple, 38
Margined-1, 32, 40
Margined-2, 32, 40
Margined-fluctuated, 35, 41
Margined-reduced, 32, 40
Margined-slight, 35
Margined-suppressed, 38
Palmate, 29, 30, 40
Pear, 35, 41
Petaloid, 39
Precocious, 28–30, 40
Purple, 32, 34, 40, 41
Rayed, 32–34, 40
Retracted, 37, 38, 41
Semi-contracted, 28, 30, 40
Shrubby, 32–34, 40
Side-reduced, 39
Smeary, 32
Speckled, 31, 35, 41
Speckled-reduced, 30, 31, 40
Star, 35
Striped, 36, 41
Terminal, 38
Tinged, 39
Tube-white, 32–34, 40
Variegated, 26–28
Variegated-reduced, 39
White-1, 35, 41
White-2, 32, 36, 41
Willow, 38
Wrinkled, 39
Yellow, 27, 30, 31, 40
Yellow-inconstant, 30
Linkage group 26
Acuminate, **32**, 35–40
Contracted, **32**, 35–40
Cordate, 27, **28**, 30–32, 35–40
Delicate, **35–41**
Duplicated, 32, **36**–39, 41
Pear, **35**, 37–39, 41
Retracted, **37**–39, 41
Speckled, **35–41**
Variegated, **26**, 28, 31, 32, 35–40
Yellow, 27, **30**–32, 35–40
Linkage studies, 26
PHIPPS, I. F., 495, 600, 610
Phylloxerans, 251
Phylogeny, 208
Physiological salt solution, 400, 405, 407–409, 413, 417

Pigeon, 397, 398, 402–412–416, **532–534**, 536, 542, **544, 547**–552, 557, 560, 563, 565, 566, 568
Albino eyes, 536, 544
Anaphylactic shock, 412
Anaphylactic state, 412, 413, 416
Anaphylaxis, 413
Breed
Barb, 533
Blondinette, 533
Carrier, 533
Dragoon, 533, 537
Homer, 533, 552, 554
Maltese, 533, **545**
Mondaine, 533
New York Flights, 533
Nun, 533
Owls, 533
Pouter, 533, **545**, 552
Runt, 533
Tumbler, 533, 553, 554
Breeding, 552
Black and chocolate flecking, **558**–562, 564, 566–568, 570
Chocolate and black flecking, **558**–562, 564, 566–568, 570
Dominant red, black and chocolate, 552, 555
Flecking, **558**–562, 564, 566–568, 570
Black, **558**–562, 564, 566–568, 570
Chocolate, **558**–562, 564, 566–568, 570
Red, 564
Relations of black, brown and red, 541
Experimental results, 537
Character
Black, 548
Chocolate, 550
Gray, 549
Intensity, 551
Color descriptions of chocolate brown types, 545, 546
Color inheritance, 548, 568
Color patterns, 548, 551
Black check, 548, 566
Black with blue tail, 548, 551
Blue barless, 548
Blue barred, 548, 565
Brown Silver, 546
Chocolate, 551
Chocolate brown, 532, 545
Dilute brown checker, 546
Dilute brown silver, 546
Dilute spread chocolate brown, 534
Self black, 548
Sooty, 548
Color variations, 547
Crossover types, 568
Gene
Extension of black or brown, 541, 543, 550
Inhibition of black, 532, 541, 544, 549, 551
Intensity of pigmentation, 532, **541**, 543, 544, 548, **551**, 552, **555**–557, 568

Pigeon (Continued)
Gene (Continued)
Non-extension of black or brown, 541, 550, 566
White, 547
Inheritance, 532, 547
Juvenile eye color and its relation to plumage color, 536
Linkage, 532, 537, 539, **541**, 543, 544, 550, 552, 555–557, 568
Black with black-inhibiting factor, 532
Brown and intensity of pigmentation, 541
Chocolate and intensity, 568
Dominant red with intensity, 555, 568
Microscopic study of pigments, 562
Mutation, 542, 567
Pigment
Black series, **562**, 568
Chocolate series, 564
Dominant red and gray, 563
Microscopic study of, 562
Pigmentation
Biochemical relationship between genes for color and pigment formation in birds and mammals, 568
Bister, 545, 546
Black, 532–**536**, **537**–543, 547, **548–552–555–558–562**, 564–566, 568, 570
Black barred, 533
Black-barred blue, 566
Black-barred silver, 534
Black, dilute clumped phase, 534
Blue, 537, 548, 549, 563, 565
Brown, 532, **533–536**, **537–539–541**, 542, 544, 565
Brown barred, 533
Brown-barred silver, 533, 537
Brown, intense-clumped phase, 534
Brown silver, 565
Buff, 545, 546
Chocolate, 534, **547**, 548, **550–552–555**, **558**, 559, 561, 562, **564**–568, 570
Chocolate brown, 532, 534, 542, **545**
Chocolate-brown dun, 534
Clumped, 549
Dilute, 551, 552, 558
Dilute black, 532, 533, 535, 536, 541, 556, 563
Dilute blue, 563
Dilute brown, 535, 536, 541, 542, 544
Dilute chocolate, 536, 555, 557, 565
Dilute red, 541, 542
Dilute spread black, 534
Distribution, 568
Dominant red, **547**, **549–552–555**, 559, 561–**563**–568, 570
Dominant yellow, 555, 564
Drab, 552, 565
Drab silver, 565
Dun, 533, 551, 558, 562, 563
Dun-barred silver, 533, 534, 537
Granules, 568
Pigeon (Continued)
Pigmentation (Continued)
Gray, **549**, 550, 558–560, **563**, 564, 566, 568, 570
Intense, 558
Intense black, 541, 544, 557, 563
Intense brown, 535, 536, 541, 542
Intense chocolate, 555–557
Intense spread brown, 534
Intense spread chocolate brown, 534
Plum color, 563
Prouts Brown, 545
Recessive red, 550, 551, 562
Red, 532, 533, 535, **536**, **537**, **539**–541, 543, **547–549–552–555**, 558–568, 570
Red barred, 565, 566
Relation of black, red and chocolate, 551
Sepia, 565
Silver, 533, 537, 563
Spread, 549
Umber, 546
Whole-colored duns, 534
Yellow, 532, 536, 544, 551
Plumage color, **532**, 542, 548, 550
Chocolate brown, **532**, 542
Juvenile eye color and its relation to, 536
Linkage of brown, 532
Relation of chocolate to black and dominant red, 547
Polypnóea, 412
Precipitating antisera, 397
Sensitized to mouse protein, anaphylactic shock in, 412
Wild Rock, 551
Pigments, chemistry of animal, 568
PINCUS, G., 44, 67, 68, 72
Plant
Hybridity, 94
Kingdom, occurrence and nature if vivipary, 587
Pollen sterility, 94
PLATE, L., 45, 52, 54
POCOCK, R. I., 50
Poisson
Law, 117, 118
Series, 114
Polyploid forms in Fragaria, **455**, 464
Polyploidy, 210
Rosa, 449
POPE, 610
Population, 110
Isolated, 114
Size, 134–137, 139–142, 146–148, 150, 151
PORTIS, B., 414
Potassium oxalate, 401
Potato, male sterility, 92
Precipitating antisera in pigeons, 397
Primula, 460
floribunda, 78, 209, 210
kewensis, 209, 473
Tetraploid, 460
verticillata, 78, 209, 210
Prunus, 479

Prunus laurocerasus, 93
Pteridophytes, 588
  Dormancy, 588
  Vivipary, 588
PUMAR, D., 52, 54
Pygaera hybrids, 209

RABAUD, E., 45, 54
Rabbit, 398, 401, 404, 406, 408–411, 413, 414, 416–418, 420
  Antiserum, 404, 416
Radium, 142, 529
Raisin-agar, 213
RANDOLPH, L. F., 473
RAPER, H. S., 567, 568
Raphanus, 450, 484
Raphanus-Brassica hybrid, 208, 209
Raphanus sativus, 484
Raspberry, 460
  La France, 460
  Tetraploid, 460
Rat, 401, 409–411, 418, 420
Rattus rattus norvegicus, **401**, 408, 411
Rattus rattus rattus, **401**, 408, 411
Rattus rattus tectorum, **401**, 408, 411
REDFIELD, H., 375, 653
REED, G. M., 303
RENNER, O., 325, 338, 339, 341, 345
Rhabdophaga heterobia, 252
Rhizophora, 589
RHOADES, M. M., 190, **375**, **490**
Rice, 291
RICHARDSON, C. W., 444, 450
RICHEY, F. D., 17
RIDGEWAY, R., 535, 544, 545
Ringer's solution, 401
ROBBINS, R. B., 100, 357–359, 363
ROBERTSON, W. R. B., **349**, 350, **353**, 355, 356
Rockefeller Institute, 161, 505
Rodent, 398, 406–409, 413, 550
  Antiserum, 398, 406, 418
  Coat color, 305
  Precipitating serum, 397, 416
  Species relationship, 410
Rodentia, 397, 410, 416
Roentgenization, 256, 262
Rooster, 414
Rosa, 75, 80, 90, 92
  acicularioides, **81**, 82
  acicularis, 77, 81, 86, 90, 91
  arkansana, 77, 79
  arvensis, 78, 83
  blanda, 77, 79, 81, 83, 85, 86, 90, 91
  Bushii, 91
  californica, 78
  carolina, 77, 83–85
  cinnamomea, 78
  Durandii, 78
  Engelmanni, 86
  Fendleri, 79
  foliolosa, 85
  humilis, 84
  Lyoni, 84
Rosa (Continued)
  microphylla, 80
  mollis, 79
  muriculata, 90
  obovata, 84
  palustris, 77, 78, 81, 85, 86, 90
  pimpinellifolia, 78
  pisocarpa, 80
  polyanthema, 89
  relicta, 79, 86, 89, 91
  rubrifolia, 92
  rudiuscula, 84, 91
  rugosa, 85, 86
  Schuettiana, **81**, 82
  sempervirens, 83
  subblanda, **83**, 85, 90
  subserrulata, 80, 81
  suffulta, 84, 91
  Tetonkaha, 85, 86
  virginiana, 77, 83, 84
  Warleyensis, 85
  Woodsii, 77
Rosaceae, 94
Roscoe B. Jackson Memorial Laboratory, 397
Rose, **75**, 80, 81, 84, 87, 89, 94
  Caninae, 78, 79, 91
  Cinnamomeae, 93
  Diploid, **80**, **85**, 89, 93, 94
  Hybrid, **75**, 80, 82, 83, **85**, 86, 89, 90, 92, 95
    Hexaploid, **86**, 90
    Sterility in interspecific, 81
    Tetraploid, **83**, 84, 86
  Hybridization, 92
  Interspecific hybrid, sterility, 81
  North American wild, 76, 93
  Polyploid, 449
  Sterility, **75**, 80, 93
    In interspecific hybrids, 81
    Ovule, 89–93
    Pollen, 75–81, 83, 84, **86**, **87**, 90–94
    Within species, 76
  Tetraploid, **80**, **83**, 84, 86, 89
  Variation in sterility within the species, 76
  Wild, **75**–79, 82, 84, 85, 87, 88, 90, 94
  Zygotic sterility, 90, 92
ROSENBERG, O., 319
Rotifers, 251
ROXAS, H. A., 560
Rubus, 78, 92, 478
  Diploid, 93
  Pollen sterility, 93
  Polyploid, 93
  rusticanus inermis, 93
  Tetraploid, 93
  thyrsiger, 93
  Triploid, 478
RYBIN, V. A., 209
Rye, 446

Saitama Agricultural Experiment Station, **1**, 22
SAX, H., 446
SAX, K., 76, 92, 93, 309, 315, **446**, **477**, **484**
SAKAMURA, T., 87, 92

SALAMAN, R. N., 92
Salix, 319
SCHEUBER, L. M., 624
SCHIEMANN, E., 444, 445
SCHMALHAUSEN, I., 222
SCHMUCK, M. L., 225
SCHRADER, F., 251
SCHRATA, 63
SCHRATZ, 72
SCHUSTER, C. E., 94
SCHWERTSCHLAGER, J., 78
Sciara, 225–229, 251, 252
  Androsomes, 227, 229, 252, 253
  Chromosome
    Dissimilarity, 250
    Male-limited, 227, 229, 252, 253
    Similarity, 249
  coprophila, 225, 229
  Crossing over, 247–249–251
  Female
    Androgenic, 228
    Gynogenic, 228
  Female-producing female, 225–228, 231, 234, 235, 239–242, 245, 246, 249, 250
  Fertility, 227, 228, 249
  Gene
    Narrow, 228–230, 231, 241, 242, 245–247, 248, 250
    Sex-linked, 227
    Sex of progeny, 225, 226, 228, 231, 232, 248–252
    Spot, 234
    Swollen, 227–230, 231, 232, 234–236, 238, 240–242, 245–247, 248, 250, 251
  Inbreeding, 227, 228
  Inheritance of narrow, 241
  Inheritance of swollen, 231, 240, 241
  Linkage of swollen and narrow, 247
  Male-producing female, 225–228, 231–234, 240, 241, 245, 246, 249, 250
  Narrow, inheritance of, 241
  Out-crosses, 227, 228, 240
  Reproduction
    Digenic, 229, 230, 238, 240
    Monogenic, 229, 240, 241
  Sex chromosome, 225–228, 249, 251, 252
  Sex determination, 225–227, 231, 251–253
  Sex-linked characters, 230, 250
  Sex-linked genes, 227
  Sex-linked inheritance of sex of progeny, 248
  Wild-type, 227–232, 234, 238, 240–242, 245, 246, 248
SCLATER, 160
Sebright bantams, 560
SEIDENBERG, 413
Selection, 98, 100, 101, 102–104, 106, 107, 111, 114, 117, 121, 127–129, 131–158
  Equilibrium, 102
  Fecundity, 101
  Genic, 103, 104
  Intensity, 135, 136
  Mating, 101, 102
  Mortality, 101
Selection (Continued)
  Natural, 146
  Pressure, 101, 103, 104, 142, 147
  Zygotic, 103, 104, 155
Self-fertilization, 357, 358, 360, 371, 373, 374
Semisterile in maize, 595
SENN, P. H., 601
Sera, antipneumococcus, 414
SEREBROVSKY, A. S., 254, 255, 264, 266
SETCHELL, W. A., 193
Sex chromosome mechanism in Sciara, 225
Sex determination in Sciara, 225
Sex-linked genes, 360, 367, 371, 373, 374
Sex-linked inheritance in Sciara, 248
SHARP, L. W., 190, 479
Sheep, 414
SHERWOOD, 414
SHREVE, R., 80, 81
Shrub, 588
  Dormancy, 588
SHULL, A. F., 251
SHULL, G. H., 99, 326, 339
Side-chain theory of the structure of the gene, 267, 276
SINGLETON, W. R., 197, 616
SIRKS, M. J., 5, 6
Size characters, 138
SLEPKOV, V. H., 254
SMITH, A. D. B., 111
Smith College, 72
SMITH, E. A., 414
SMITH, W. K., 300
SNELL, G. D., 42, 60
SO, M., 47
SOBOTKA, 414
Solanum, 93, 446
  Diploid, 93
  Melongena (See eggplant), 1
  nigrum, 479
  Pollen sterility, 93
  Tetraploid, 93
  tuberosum, 87, 588
SOLMS-LAUBACH, H., 441
SORGHUM, 291–295, 298, 299, 301, 302, 304–307
  Albinism, 302
  Albino seedings, 293, 296, 300, 302–305, 307
  Blackhul kafir, 294
  Chlorophyll, 291–307
    Deficiencies, 299, 300, 304–307
    Mutations in, 306, 307
    Seedling characters, 291–307
  Green seedlings, 293–305, 307
  Hybrid vigor, 302
  Inheritance, 291, 293, 303–305
    Of chlorophyll characters, 291, 293, 304–306
  Kafir, 291–294, 297, 299, 301, 302, 307
    Chlorophyll deficiencies, 292–294, 299
    Inheritance, 297, 301
    Mutation, 293, 297, 298, 302
  Linkage, 291, 292, 302, 304, 307
    Red stem color and albinism, 302
    Relations, 291

Sorghum (Continued)
Linkage (Continued)
Relations of chlorophyll characters, 291, 292
Red stem albino, 302–304
Red stem color, **302**–304, 307
Striped green seedlings, 305, 307
Striped yellow seedlings, 305, 307
Sudan grass, 300–305, 307
Virescent seedlings, **294**–299, 301, 307
White seedlings, **292**–294, 303, 304, 307
Yellow seedlings, **300**, 301, 305, 307
Spencer apochromatic objective, 424, 474
Spermatophytes, 588
Dormancy, 588
Vivipary, 588
STADLER, L. J., 99, 306, 529
Staining, gentian-violet-KKI-clove oil method, 645
STEELE, D. G., **532**, 537, 550–552, 558, 559
STEERE, W. C., 347
Step allelomorphism in Drosophila, **254**, 255, 262–265
Sterility in wild roses and in some species hybrids, 75
STERN, C., 393, 630
Stizolobium, 92, 595, 597
Storrs Agricultural Experiment Station, 532, 533, 538, 542, 544
STOW, I., 87, 92
STROMAN, G. N., 300
STURTEVANT, A. H., 57, **105**, 269–271, 273–276, 279, 280, 346, 375–377, 379–381, 389, 393, 491, 630, 636, 644
Sudan grass (See sorghum), 300–305, 307
SUMNER, F. B., 165
SUTTON, I., 94

Tailrings, cross between mice with different numbers of, 591
TATEISI, T., 17
TEGETMEIER, 537
Temperature, 142
TENBROECK, C., 161
Tetraploidy, 144
Tettigidae, 350, 353–355
Inbreeding, 356
Origin of partheno-produced males, **353**–356
parvipennis, 355
Texas Agricultural Experiment Station, **291**, 455
THIGPEN, 66, 72
THOMPSON, D. H., **267**, 271
THOMPSON, W. P., 92, 196, **309**, 312, 315, 440, 446, 463, 485
TICE, S. C., 268
Timiriaziev Biological Institute, 254
TISCHLER, G., 478
Tokyo Imperial University, 26
Tomato, 14, 17, 484
Dwarf Champion, 14
Stone, 14
Translocation, 144
In Drosophila, 490
Involving the second and fourth chromosomes of Drosophila, **629**–632, 653
Tree, dormancy, 588
Triticum, 92, 93, 309, 446
dicoccoides, 310, 447
dicoccum, 310
Diploid, 93
durum, 310–312, 318, 319, 321, 322, 447, 484
Hexaploid, 93
Hybrid, 446
monococum, 309–312, 314–316, 318, 319, 321–323
Pentaploid, 93
polonicum, 310, 447
Tetraploid, 93
Triploid, 93
turgidum, 309, 310, 312, 322, 446
vulgare, 196, 309, 312, 314, 315, 320, 322, **374**, 446, 447, 485
vulgare-durum, 196
Troemmer diluting pipette, 416
TROMMSDORF, R., 410, 413, 416, 418
TSCHERMAK, E., 5
Tuberculin luer, 399
Tulip, triploid, 477
Tulipa Clausiana, 445
Tyrosinase, 567, 568
Tyrosine, 567, 568

UHLENHUTH, 410, 416, 418
ULLIAN, S. S., 225, 227, 230, 231, 234
United States Department of Agriculture, 624
United States Weather Bureau, 436
University of
California, **191**, 323
Chicago, **97**, 659, 669
Chicago Hospital, 659, 669
Illinois, 267, 288, 289, 612
Latvia, 43, 52
Michigan, **75**, 76, 85, 325, 386, 387, **397**, 398
Michigan Hospital, 401
Minnesota, 624
Missouri, 574
Pittsburgh, 505, 659
Saskatchewan, **309**, 323
Wisconsin, 532, 533, 545, 547, 548, **595**, 624

VAN FLEET, W., **444**
Variability of characters, 139
Variation, 97, **106**
Gene frequency, 100, 106
Random, 151
Viability, effect of X-ray radiation on Habrobracon, **505**, **527**, 528
Victoreen Dosimeter, 661
Viola, 450
Vitis, 93
Hybrid, 93
Pollen sterility, 93
vulpina, 93

Vivipary, **574, 575–578–581**–584, 586, **587**–590
Eucalptodon, 588
Fegatella, 588
Maize, **574, 575–578–581**–584, 586, **587**, 589, 590
Occurrence and nature of in the plant kingdom, **587**, 590
VRIES, DE, H., 98, 99

WADDINGTON, C. H., 357
WADE, B. L., 624
WAGNER, 98
WAHLUND, S., 128
WALDO, G. F., 437
WALL, C. N., 289
Wasp (See Habrobracon), 280, **505**, 530, 659, 661
WATKINS, A. E., 76, 94
WEIDANZ, 410, 416, 418
WEINBERG, W., 100
WEINSTEIN, A., 490, 528
WELLINGTON, 17
Western Pennsylvania Hospital, 416
Wheat, 5, 76, 94, 291, 300, **309**–311, 315, 319, 323, 446, 484
Backcrossing, 311, 316
Cross, **309**, 323
Cross-pollination, 311, 314, 320
Cytology, 309
Cytology and genetics of crosses between fourteen- and seven-chromosome species of, 309
Genetics, 309
Hybrid, 94, **484**, 485
Inheritance, 321, 322
Out-crossing, **314**, 321, 322
Pentaploid hybrid, **484**, 485
Pollen abortion, 321
Self-pollination, 311
Species
7-chromosome, **309**–311
12-chromosome, 315
13-chromosome, 315, 323
14-chromosome, **309**–311, 314–317, 319, 320, 323
15-chromosome, **314**, 315
16-chromosome, 315
21-chromosome, 314, 315, 317, 318, 323
26-chromosome, 320, 323
28-chromosome, 312, 313, 320, 321, 323
29-chromosome, 314, 320, 323
35-chromosome, 314, 320, 323
Wheat (Continued)
Species (Continued)
36-chromosome, **314**, 321
42-chromosome, 312, 314, 315, 321, 323
Species cross, **309, 311, 315–321**, 323
WHITING, A. R., 280, 506, 512, 513, 529, **659**, 667
WHITING, P. W., 505, 506, 510, 513, 516, 520, 525, 529, 659, 660, 666, 667
WILLMOTT, E. A., 85
WILLSTÄTTER, R., 287
WILLSTÄTTER's structure of enzymes, 287
WILSON, E. B., 229, 251
WINGARD, S. A., 6
Wisconsin Agricultural Experiment Station, 600, 601
WOLFE, T. K., 5
WOODWORTH, C. M., 612
WRIEDT, C., 536, 537, 539, 544, 548, 550, 552, 557–561
WRIGHT, S., 46, **97**, 101, 103, 106, 107, 109–111, 129, 133, 152, 279, 280, 368, 373, 562, 565, 566

X-ray, 45, 54, 55, 92, 99, 142, 254, 255, 306, 346, 380, 479, 480, 490, 491, 498, 630, 653, 659–661, 669, 670, 678, 679
Radiation on Habrobracon, 505, 507, 526, 527
Wappler Diex, 506
Xanthophyll, 299

Yale University, 533, 538, 542, 544
YARNELL, S. H., **422, 455**, 461, 464, 472, 478
YASUI, K., 39
Yeast, 213, 228
YULE, G. U., 508, 615, 616

Zea mays, (See maize) 92, **175**–177, 181, 186, 190, 473, 479, 484
Cytological identification of chromosome associated with *r-g* linkage group, 175
Inheritance, disomic, 178, 187–190
Inheritance, trisomic, 175, **176**, 178, 181, 186, 187, 190
Linkage groups, **175**, 178–180, 185, **186**, 187, 189, 190
Triploid, 484
ZEIMET, A., 545
Zeiss objective, 646
ZELENY, C., 267, 269, 270, 276–278, 288
ZINSSER, H., 412